Changes in *Helicobacter pylori* Immunoglobulin G Levels and Gastric Mucosal Atrophy after Successful Eradication of *Helicobacter pylori*

Ji Hye Kim^{1,*}, Joon Seop Lee^{1,*}, Yong Hwan Kwon^{1,2}, Seong Woo Jeon^{1,2}, Su Youn Nam¹, Sun Jin¹

Department of Internal Medicine, Kyungpook National University Hospital¹, Department of Internal Medicine, School of Medicine, Kyungpook National University², Daegu, Korea

Korean J *Helicobacter* Up Gastrointest Res 2019;19:174-183 <https://doi.org/10.7704/kjhugr.2019.19.3.174>

대한상부위장관·헬리코박터학회지에 게재된 위 논문에서 Fig. 1에 오류가 있어 정정합니다.

This correction is being published to correct the Fig. 1 in above article.

Before correction

Fig. 1. Schematic flow of this study. ESD, endoscopic submucosal dissection; EGC, early gastric cancer; *H. pylori*, *Helicobacter pylori*; IgG, immunoglobulin G; PG, pepsinogen.

Corresponding author: Yong Hwan Kwon

Department of Internal Medicine, Kyungpook National University Hospital, 130 Dongdeok-ro, Jung-gu, Daegu 41944, Korea

Tel: +82-53-200-2609, Fax: +82-53-200-3089, E-mail: tear9754006@yahoo.co.kr

*These authors contributed equally to this work.

After correction

Fig. 1. Schematic flow of this study. ESD, endoscopic submucosal dissection; EGC, early gastric cancer; *H. pylori*, *Helicobacter pylori*; IgG, immunoglobulin G; PG, pepsinogen.